

**PATIENT INFORMATION**  
**PROAIR RESPICLICK (prō'ār res-pē-kliik)**  
**(albuterol sulfate)**  
**Inhalation Powder**

**What is PROAIR RESPICLICK?**

PROAIR RESPICLICK is a prescription medicine used in people 4 years of age and older to:

- treat or prevent bronchospasm in people who have reversible obstructive airway disease
- prevent exercise-induced bronchospasm

It is not known if PROAIR RESPICLICK is safe and effective in children under 4 years of age.

**Who should not use PROAIR RESPICLICK?**

**Do not use PROAIR RESPICLICK if you** are allergic to albuterol sulfate, lactose, milk proteins, or any of the ingredients in PROAIR RESPICLICK. See the end of this leaflet for a complete list of ingredients in PROAIR RESPICLICK.

**What should I tell my doctor before using PROAIR RESPICLICK?**

**Before you use PROAIR RESPICLICK, tell your doctor if you:**

- have heart problems
- have high blood pressure (hypertension)
- have convulsions (seizures)
- have thyroid problems
- have diabetes
- have low potassium levels in your blood
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if PROAIR RESPICLICK will harm your unborn baby. Talk to your doctor if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if PROAIR RESPICLICK passes into your breast milk. Talk to your doctor about the best way to feed your baby if you are using PROAIR RESPICLICK.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. PROAIR RESPICLICK and other medicines may affect each other and cause side effects. PROAIR RESPICLICK may affect the way other medicines work, and other medicines may affect the way PROAIR RESPICLICK works.

Especially tell your doctor if you take:

- other inhaled medicines
- beta blocker medicines
- diuretics
- digoxin
- monoamine oxidase inhibitors
- tricyclic antidepressants

Ask your doctor or pharmacist for a list of these medicines if you are not sure.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

**How should I use PROAIR RESPICLICK?**

- For detailed instructions, see "Instructions for Use" at the end of this Patient Information.
- Use PROAIR RESPICLICK exactly as your doctor tells you to use it.
- If your child needs to use PROAIR RESPICLICK, watch your child closely to make sure your child uses the inhaler correctly. Your doctor will show you how your child should use PROAIR RESPICLICK.
- Each dose of PROAIR RESPICLICK should last up to 4 hours to 6 hours.
- **Do not** increase your dose or take extra doses of PROAIR RESPICLICK without first talking to your doctor.
- Do not use a spacer or volume holding chamber with PROAIR RESPICLICK. PROAIR RESPICLICK does not need priming.
- Get medical help right away if PROAIR RESPICLICK no longer helps your symptoms.
- Get medical help right away if your symptoms get worse or if you need to use your inhaler more often.
- While you are using PROAIR RESPICLICK, **do not** use other inhaled rescue medicines and asthma medicines unless your doctor tells you to do so.

**PROAIR RESPICLICK (albuterol sulfate) Inhalation Powder**

- Call your doctor if your asthma symptoms like wheezing and trouble breathing become worse over a few hours or days. Your doctor may need to give you another medicine (for example, corticosteroids) to treat your symptoms.

**What are the possible side effects of PROAIR RESPICLICK?**

**PROAIR RESPICLICK may cause serious side effects, including:**

- **worsening trouble breathing, coughing and wheezing (paradoxical bronchospasm).** If this happens stop using PROAIR RESPICLICK and call your doctor or get emergency help right away. Paradoxical bronchospasm is more likely to happen with your first use of a new asthma inhalation medicine.
- **heart problems, including faster heart rate and higher blood pressure**
- **possible death in people with asthma who use too much PROAIR RESPICLICK**
- **allergic reactions.** Call your doctor right away if you have the following symptoms of an allergic reaction:
  - itchy skin
  - rash
  - swelling beneath your skin
  - worsening trouble breathing or in your throat
- **low potassium levels in your blood**
- **worsening of other medical problems in people who also use PROAIR RESPICLICK including increases in blood sugar**

The most common side effects of PROAIR RESPICLICK include:

- back pain
- body aches and pain
- upset stomach
- sinus headache
- urinary tract infection
- your heart feels like it is pounding or racing (palpitations)
- chest pain
- fast heart rate
- shakiness
- nervousness
- headache
- dizziness
- sore throat
- runny nose

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of PROAIR RESPICLICK. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store PROAIR RESPICLICK?**

- Store PROAIR RESPICLICK at room temperature between 59°F and 77°F (15°C and 25°C).
- Avoid exposure to extreme heat, cold, or humidity.
- Keep the cap on the inhaler closed during storage.
- Keep your PROAIR RESPICLICK inhaler dry and clean at all times.

**Keep PROAIR RESPICLICK and all medicines out of the reach of children.**

**General Information about the safe and effective use of PROAIR RESPICLICK**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use PROAIR RESPICLICK for a condition for which it was not prescribed. Do not give PROAIR RESPICLICK to other people, even if they have the same symptoms that you have. It may harm them.

This Patient Information leaflet summarizes the most important information about PROAIR RESPICLICK. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about PROAIR RESPICLICK that is written for health professionals. For more information, go to [www.MyProAir.com](http://www.MyProAir.com) or call 1-888-482-9522.

**What are the ingredients in PROAIR RESPICLICK?**

Active ingredient: albuterol sulfate

Inactive ingredients: lactose (may contain milk proteins)

Marketed by: Teva Respiratory, LLC, Horsham, PA 19044

Manufactured by: Teva Pharmaceutical Industries, Ltd. Jerusalem, Israel

Copyright ©2015-2016, Teva Respiratory, LLC All rights reserved  
ProAir® and RespiClick® are trademarks owned by Teva Respiratory, LLC  
PRORPL-003 Rev. 09/16

## PROAIR RESPICLICK (albuterol sulfate) Inhalation Powder

### Instructions for Use PROAIR RESPICLICK (prō'ār res-pē-klik) (albuterol sulfate) inhalation powder

#### Your PROAIR RESPICLICK Inhaler

When you are ready to use PROAIR RESPICLICK for the first time, remove the PROAIR RESPICLICK inhaler from the foil pouch.

There are 2 main parts of your PROAIR RESPICLICK inhaler including:

- the white inhaler with the mouthpiece. **See Figure A.**
  - the red cap that covers the mouthpiece of the inhaler. **See Figure A.**
- There is a dose counter in the back of the inhaler with a viewing window that shows you how many doses of medicine you have left. **See Figure A.**




Figure A

- Your PROAIR RESPICLICK inhaler contains 200 doses (inhalations). **See Figure B.**
- The dose counter shows the number of doses left in your inhaler.
- When there are 20 doses left, the dose counter will change to red, and you should refill your prescription or ask your doctor for another prescription.
- When the dose counter displays '0,' your inhaler is empty, and you should stop using the inhaler and throw it away. **See Figure B.**



Figure B

#### IMPORTANT:

- **Always close the cap after each inhalation so your inhaler will be ready for you to take your next dose.** Do not open the cap unless you are ready for your next dose.
- You will hear a “click” sound when the cap is opened fully. If you do not hear the “click” sound the inhaler may not be activated to give you a dose of medicine.
- **PROAIR RESPICLICK does not have an activation button or medicine canister.** When you open the cap, a dose of PROAIR will be activated for delivery of the medicine.
- In general, the technique for administering PROAIR RESPICLICK to children is similar to that for adults. Children should use PROAIR RESPICLICK under adult supervision, as instructed by the patient’s physician.
- Do not use a spacer or volume holding chamber with PROAIR RESPICLICK. PROAIR RESPICLICK does not need priming.

#### Using your PROAIR RESPICLICK inhaler:

**Important: Make sure the red cap is closed before you start using your inhaler.**

## PROAIR RESPICLICK (albuterol sulfate) Inhalation Powder

### Step 1. Open

- Hold the inhaler upright and open the red cap fully until you feel and hear a “click”. **See Figure C.**
- Each time you open the red cap and it “clicks”, a dose of PROAIR RESPICLICK is ready to be inhaled.




Figure C

#### Remember:

- For the correct use of PROAIR RESPICLICK, **hold the inhaler upright** as you open the red cap. **See Figure D.**
- **Do not** hold the inhaler in any other way as you open the red cap.
- **Do not** open the red cap until you are ready to take a dose of PROAIR RESPICLICK.




Figure D

### Step 2. Inhale

- Before you inhale, breathe out (exhale) through your mouth and push as much air from your lungs as you can. **See Figure E.**
- **Do not** exhale into the inhaler mouthpiece.




Figure E

- Put the mouthpiece in your mouth and close your lips tightly around it. **See Figure F.**




Figure F

## PROAIR RESPICLICK (albuterol sulfate) Inhalation Powder

- **Do not** block the vent above the mouthpiece with your lips or fingers. **See Figure G.**




Figure G

- Breathe in quickly and deeply through your mouth, to deliver the dose of medicine to your lungs.
- Remove the inhaler from your mouth.
- Hold your breath for about 10 seconds or for as long as you comfortably can.
- Your PROAIR RESPICLICK inhaler delivers your dose of medicine as a very fine powder that you may or may not taste or feel. **Do not** take an extra dose from the inhaler even if you do not taste or feel the medicine.

### Step 3. Close




Figure H

- Close the red cap firmly over the mouthpiece. **See Figure H.**
- Make sure you close the red cap after each inhalation so that the inhaler will be ready for your next dose.
- **If you need another dose** close the red cap and then repeat steps 1-3.



### How should I store PROAIR RESPICLICK?

- Store PROAIR RESPICLICK at room temperature between 59°F and 77°F (15°C and 25°C).
- Avoid exposure to extreme heat, cold, or humidity.
- Keep the red cap on the inhaler closed during storage.
- Keep your PROAIR RESPICLICK inhaler dry and clean at all times.
- **Keep your PROAIR RESPICLICK inhaler and all medicines out of the reach of children.**

### Cleaning your PROAIR RESPICLICK inhaler

- **Do not wash or put any part of your PROAIR RESPICLICK inhaler in water.**
- PROAIR RESPICLICK contains a powder and must be kept clean and dry at all times.
- If the mouthpiece needs cleaning, gently wipe it with a dry cloth or tissue.

## PROAIR RESPICLICK (albuterol sulfate) Inhalation Powder

### Replacing your PROAIR RESPICLICK inhaler

- The counter on the back of your inhaler shows how many doses you have left.
- When there are 20 doses left, the dose counter color will change to red, and you should refill your prescription or ask your doctor for another prescription.
- When the counter displays '0' your PROAIR RESPICLICK inhaler is empty, and you should stop using the inhaler and throw it away.
- Throw away PROAIR RESPICLICK 13 months after removing it from the foil pouch for the first time, when the dose counter displays '0', or after the expiration date on the package, whichever comes first.

### Important information

- Do not open the red cap unless you are taking a dose. Repeatedly opening and closing the cap without inhaling a dose will waste the medicine and may damage your inhaler.
- Your PROAIR RESPICLICK inhaler contains dry powder so it is important that you do not blow or breathe into it.

### Support

- If you have any questions about PROAIR RESPICLICK or how to use your inhaler, go to [www.ProAir.com](http://www.ProAir.com) or call 1-888-482-9522.

This Patient Information and Instructions for Use have been approved by the U.S. Food and Drug Administration.

Marketed by:

Teva Respiratory, LLC  
Horsham, PA 19044

Manufactured by:

Teva Pharmaceutical Industries, Ltd.  
Jerusalem, Israel

Copyright ©2015-2016, Teva Respiratory, LLC All rights reserved

ProAir® and RespiClick® are registered trademarks owned by Teva Respiratory, LLC.



PRORIFU-003 Rev. 09/2016

PRS-40781